



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2021

---

## **Observation of Collagen-Containing Lesions After Hematoma Resolution in Intracerebral Hemorrhage**

Love, Christopher J ; Kirschenbaum, Daniel ; Selim, Magdy ; Lo, Eng H ; Rushing, Elisabeth ; Spector, Myron ; Aguzzi, Adriano

**Abstract:** Background and purpose: The classic presentation of chronic (stage III) hemorrhagic stroke lesions is a fluid-filled cavity. In one of the most commonly used animal models of intracerebral hemorrhage (ICH), we noticed additional solid material within the chronic lesion. We examined the composition of those chronic ICH lesions and compared them with human autopsy cases. Methods: ICH was induced in rats by the injection of collagenase in the striatum. Tissue sections after hematoma resolution corresponding to 3 different chronic time points-28, 42, and 73 to 85 days post-ICH-were selected. Human autopsy reports at the University Hospital of Zurich were searched between 1990 and 2019 for ICH, and 3 chronic cases were found. The rat and human sections were stained with a variety of histopathologic markers. Results: Extensive collagenous material was observed in the chronic lesion after hematoma resolution in both the rat model and human autopsy cases. Additional immunostaining revealed that the material consisted primarily of a loose network of collagen 3 intermingled with occasional GFAP (glial fibrillary acidic protein)-positive processes and collagen 4. Conclusions: A key feature of the chronic ICH lesion is a loose network of collagen 3. The collagenase rat model reproduces the morphology and composition of the chronic human ICH lesion. While identifying new features of ICH lesion pathology, these results are important for treatment and recovery strategies.

DOI: <https://doi.org/10.1161/strokeaha.120.030240>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-201619>

Journal Article

Accepted Version

Originally published at:

Love, Christopher J; Kirschenbaum, Daniel; Selim, Magdy; Lo, Eng H; Rushing, Elisabeth; Spector, Myron; Aguzzi, Adriano (2021). Observation of Collagen-Containing Lesions After Hematoma Resolution in Intracerebral Hemorrhage. *Stroke*, 52(5):1856-1860.

DOI: <https://doi.org/10.1161/strokeaha.120.030240>

**Observation of collagen-containing lesions after hematoma resolution in  
intracerebral hemorrhage**

Christopher J. Love, PhD\*; Daniel Kirschenbaum, MD, PhD\*; Magdy Selim, MD, PhD;  
Eng H. Lo, PhD; Elisabeth Rushing, MD; Myron Spector, PhD; and Adriano Aguzzi,  
MD, PhD

\*equal contribution

**Affiliations**

Christopher J. Love:

Department of Mechanical Engineering, Massachusetts Institute of Technology.

Daniel Kirschenbaum:

Institute of Neuropathology, University of Zurich.

Magdy Selim:

Department of Neurology, Beth Israel Deaconess Medical Center, Harvard  
Medical School.

Eng H. Lo:

Departments of Radiology and Neurology, Massachusetts General Hospital,  
Harvard Medical School.

Elisabeth Rushing:

Institute of Neuropathology, University of Zurich.

Myron Spector:

Department of Orthopedics, Brigham and Women's Hospital, Harvard Medical School; Tissue Engineering and Regenerative Medicine, VA Boston Healthcare System.

Adriano Aguzzi:

Institute of Neuropathology, University of Zurich.

### **Corresponding author**

Adriano Aguzzi

Institute of Neuropathology, University of Zurich

Schmelzbergstrasse 12

8091 Zurich

[adriano.aguzzi@usz.ch](mailto:adriano.aguzzi@usz.ch)

+41 44 255 20 18

**Keywords:** stroke, intracerebral hemorrhage, hydrogels, neuropathology, collagen

**Subject terms:** Intracranial Hemorrhage, Cerebrovascular Disease/Stroke, Basic Science Research, Animal Models of Human Disease, Translational Studies

**Tables 0; Figures 2**

**Total Word Count: 2421**

**Background and purpose**—The classic presentation of chronic (Stage III) hemorrhagic stroke lesions is a fluid-filled cavity. In one of the most commonly used animal models of intracerebral hemorrhage (ICH), we noticed additional solid material within the chronic lesion. We examined the composition of those chronic ICH lesions and compared them with human autopsy cases.

**Methods**—ICH was induced in rats by the injection of collagenase in the striatum. Tissue sections after hematoma resolution corresponding to three different chronic time points—28, 42, and 73-85 days post-ICH—were selected. Human autopsy reports at the University Hospital of Zurich were searched between 1990 and 2019 for ICH and three chronic cases were found. The rat and human sections were stained with a variety of histopathological markers.

**Results**—Extensive collagenous material was observed in the chronic lesion after hematoma resolution in both the rat model and human autopsy cases. Additional immunostaining revealed that the material consisted primarily of a loose network of collagen 3 intermingled with occasional GFAP-positive processes and collagen 4.

**Conclusions**—A key feature of the chronic ICH lesion is a loose network of collagen 3. The collagenase rat model reproduces the morphology and composition of the chronic human ICH lesion. While identifying new features of ICH lesion pathology, these results are important for treatment and recovery strategies.

## **Nonstandard Abbreviations and Acronyms**

ICH    intracerebral hemorrhage

GFAP    glial fibrillary acidic protein

NeuN neuronal nuclear protein  
 $\alpha$ -SMA alpha smooth muscle actin  
H&E hematoxylin and eosin  
eVG elastic Van Gieson

## **Introduction**

The evolution of hemorrhagic stroke lesions consists of three stages: deformation, edema, and necrosis of the surrounding tissues (Stage I); resorption of the hematoma (Stage II); and full resolution of the hematoma resulting in a fluid-filled cavity (Stage III).<sup>1</sup> In human Stage II lesions, a capsule-like, collagenous border has been observed.<sup>2,3</sup> Histological examination after autologous blood injection in dogs reveals the formation of argentophilic fibers on the border of the hematoma with a gradual change to more mature collagenous fibers. However, the studies were limited to Stages I and II with the hematoma still present.<sup>4,5</sup> Prior reports of Stage III hemorrhagic lesions in humans describe a thin, fluid-filled cavity that is indistinguishable from a Stage III infarct except for the presence of blood-related deposits such as hemosiderin and walls and fluid that have a yellow to brown coloration.<sup>1,2,6,7</sup>

Histopathological characterization of chronic stroke lesions is important for the development of new biomaterial-based therapies<sup>8</sup> that physically interact with lesion constituents and require a space to accommodate the injection. Development of these emerging therapies typically begins in rodent stroke models, but lesion outcomes may be species dependent.<sup>9</sup> Therefore, histopathological comparison is essential to determine if chronic lesions in animal models reproduce the key features found in clinical cases.

While developing a biomaterial-based therapy<sup>10</sup> for chronic ICH in a collagenase-induced rat model<sup>11</sup>, we noticed that the chronic rat lesions were not a simple fluid-filled cavity as expected. Herein, we report on the composition of these lesions and compare them with human autopsy cases of chronic ICH.

## Methods

This study adheres to the AHA journals' implementation of the Transparency and Openness Promotion Guidelines; the authors declare that all supporting data are available within the article (and in the Data Supplement). Human autopsy reports at the University Hospital of Zurich were searched between 1990 and 2019 for ICH and three chronic cases (1 male and 2 females) were found. The Executive Director of the Kantonale Ethikkommission provided notification that the histopathologic examination of autopsy samples did not fall within the scope of the Human Research Act (HRA). Human samples consisted of formalin-fixed tissue embedded in paraffin with immunostaining for GFAP, collagen 3, collagen 4, NeuN, and  $\alpha$ -SMA by standard methods based on the oxidation of 3,3'-diaminobenzidine.

The rat model consisted of injection of collagenase (0.1U type VII in 1 $\mu$ L saline) into the right striatum of male Sprague-Dawley rats (280-300 grams). All procedures were performed in accordance with the Institutional Animal Care and Use Committee of Massachusetts General Hospital (animal protocol number 2012N000165) and the Guide for Use and Care of Laboratory Animals from the National Institute of Health (NIH). At 28, 42, and 73-85 days post-ICH, the rats (n=6 per group) were sacrificed and the extracted brains were fixed in 4% paraformaldehyde, cryopreserved in 30% sucrose, and

frozen. Cryosections (30 $\mu$ m) were immunostained for GFAP, collagen 3, collagen 4, NeuN, and  $\alpha$ -SMA by standard methods based on a secondary fluorescent antibody.

Additional experimental details for the human and rat samples are provided in the Data Supplement.

## Results

In a well-validated, collagenase-induced rat model of ICH<sup>11</sup>, we noticed an unexpected material inside chronic lesions after hematoma resolution (Figure 1). Further staining revealed that the material consisted primarily of collagen 3 intermingled with occasional GFAP-positive processes and collagen 4. Collagen 3 is typically associated with reticular fibers and forms the loose extracellular matrix found early within a wound that heals by coagulative necrosis. There was no significant difference over time in the amount of collagen 3 or lesion volume that would indicate wound healing (Figure I in the Data Supplement). The formation of a stable, hemorrhagic lesion in rat after hematoma resolution (~ 2-3 weeks depending on initial hematoma size) is consistent with prior studies.<sup>12,13</sup>

At all three examined time points and for reasons that are not yet known, we observed two types of lesion morphology: thin, compacted lesions and larger, less dense lesions. The collagenous fibers were found within both lesion types (Figure 1). NeuN (Figure II in the Data Supplement) and  $\alpha$ -SMA (Figure III in the Data Supplement) were absent within the lesions at all time points. Negative controls are included in the Data Supplement (Figure IV).

To check our rat model observations with human outcomes, we searched autopsy reports at the University Hospital of Zurich between 1990 and 2019 for ICH and found three Stage III cases (Table and Figure V in the Data Supplement). Stage III hemorrhagic lesions were not the direct cause of death in any of the patients.

An unexpected finding was the extensive extracellular matrix (ECM) inside the lesion in the H&E stained sections (Figure 2). Subsequent elastic Van Gieson (eVG) staining (Figure 2) revealed that most of the material consisted of loosely organized collagen fibers (in pink). We performed additional immunostaining, which revealed predominantly type 3 collagen (Figure 2). GFAP, non-vascular  $\alpha$ -SMA, and NeuN were negative within the lesions (Figure VI in the Data Supplement). The lack of pervasive  $\alpha$ -SMA is consistent with the finding of a loose network of collagen 3 in contrast to the contracted collagen 1 scar of chronic skin and musculoskeletal lesions. Some calcification was observed in the lesion of patient 2. Negative controls are included in the Data Supplement (Figure VII).

## **Discussion**

We observed extensive collagenous material, primarily collagen 3 with occasional GFAP-positive processes and collagen 4, in the chronic lesion after hematoma resolution in both the rat model and human autopsy cases. The source of the collagen-producing cells is unknown. Proposed sources based on brain and spinal cord lesions in rodents include meningeal fibroblasts<sup>14</sup>, perivascular fibroblasts<sup>15</sup>, and endothelial cells<sup>16</sup>. The proposed purpose of the collagen in rodent lesions is to provide a scaffold for invading fibroblasts, macrophages, and blood vessels.<sup>14</sup> In Stage II ICH human lesions, we observe a collagen capsule surrounding the hematoma that is distinct



from an outer GFAP-positive glial layer, which contains CD68-positive microglia/macrophages (Figure VIII in the Data Supplement). This subacute capsule consists primarily of collagen 3 (Figure VIII in the Data Supplement); therefore, the collagen 3 observed in Stage III lesions might be related to the Stage II capsule.

We were surprised to find only a few chronic cases of ICH in the archives. Perhaps chronic ICH lesions are underrepresented because incidence is lower and acute mortality is higher compared to ischemic stroke.<sup>17</sup> Nowadays, very few autopsies for ICH are performed because they can be diagnosed by neuroimaging. While old microbleeds are difficult to detect, we tried looking at cavernous hemangiomas, which are known to bleed repeatedly with a prolonged history; however, the distinction between collagen associated with atypical vessels and putative bleeding-associated tissue is difficult. Further studies of larger numbers of patients would be helpful to confirm our findings and expand our understanding of them—such as if they are global or restricted to ICH in specific brain regions.

Despite the very different time of investigation post-ICH of the rat (tens of days) and human Stage III lesions (years), they are found with the same morphology and composition. Our comparison is based on the end stage of the lesion, and further work is needed to determine if these findings are coincidental or related to similarities in hematoma resolution with a different time scale based on the initial hematoma size.

The examination and comparison of chronic stroke lesions in animal models and human clinical cases is important for treatment and recovery strategies. The injection of a biomaterial-based therapy into a chronic lesion requires space to accommodate the biomaterial. The loose, porous collagen 3 matrix that was observed unexpectedly in the

chronic rat lesion did not inhibit the subsequent injection of a biomaterial into the lesion.<sup>10</sup> Presumably, the liquid injection displaces the collagen fibers or infiltrates the porous network before gelation in situ. Based on our findings of a similar lesion composition in the human cases, we anticipate that chronic human ICH lesions can also accommodate and prove amenable to treatment with a biomaterial-based therapy.

**Sources of Funding.** The authors gratefully acknowledge a travel grant from the MIT MISTI program and support from the AHA Collaborative Sciences Award (15CSA24540001), Swiss National Research Foundation, and Nomis Foundation.

**Disclosures.** None.

### **Supplemental Materials**

Table

Figures I - IX

Expanded Materials and Methods

### **References**

1. Case C, Kaplan L. Intracerebral Hemorrhage. Butterworth-Heinemann; 1994.
2. Stehbens WE. Pathology of the cerebral blood vessels. C. V. Mosby; 1972.
3. Lee Y, Moser R, Bruner J, Van Tassel P. Organized intracerebral hematoma with acute hemorrhage: CT patterns and pathologic correlations. *American Journal of Roentgenology*. 1986;147:111–118.

4. Takasugi S, Ueda S, Matsumoto K. Chronological changes in spontaneous intracerebral hematoma--an experimental and clinical study. *Stroke*. 1985;16:651–658.
5. Enzmann DR, Britt RH, Lyons BE, Buxton JL, Wilson DA. Natural History of Experimental Intracerebral Hemorrhage: Sonography, Computed Tomography and Neuropathology. *AJNR*. 1981;517–526.
6. Courville C. Pathology of the Central Nervous System: A Study Based Upon a Survey of Lesions Found in a Series of Forty Thousand Autopsies. 3rd ed. Pacific Press Publishing Association; 1950.
7. Love S, Greenfield JG. Greenfield's Neuropathology. 9th ed. Boca Raton: CRC Press; 2015.
8. Love CJ, Selim M, Spector M, Lo EH. Biomaterials for Stroke Therapy: Key Questions in Research and Clinical Translation. *Stroke*. 2019;50:2278–2284.
9. Inman DM, Steward O. Physical Size Does Not Determine the Unique Histopathological Response Seen in the Injured Mouse Spinal Cord. *Journal of Neurotrauma*. 2003;20:33–42.
10. Lim TC, Mandeville E, Weng D, Wang L-S, Kurisawa M, Leite-Morris K, et al. Hydrogel-Based Therapy for Brain Repair After Intracerebral Hemorrhage. *Translational Stroke Research*. 2019.


11. Rosenberg GA, Mun-Bryce S, Wesley M, Kornfeld M. Collagenase-induced intracerebral hemorrhage in rats. *Stroke*. 1990;21:801–807.
12. Marc R. Del Bigio, Hui-Jin Yan, Richard Buist, and James Peeling. Experimental Intracerebral Hemorrhage in Rats: Magnetic Resonance Imaging and Histopathological Correlates. *Stroke*. 1996;27:2312–2320.
13. MacLellan CL, Silasi G, Poon CC, Edmundson CL, Buist R, Peeling J, Colbourne F. Intracerebral Hemorrhage Models in Rat: Comparing Collagenase to Blood Infusion. *J Cereb Blood Flow Metab*. 2008;28:516–525.
14. Maxwell WL, Duance VC, Lehto M, Ashurst DE, Berry M. The distribution of types I, III, IV and V collagens in penetrant lesions of the central nervous system of the rat. *Histochem J*. 1984;16:1219–1229.
15. Soderblom C, Luo X, Blumenthal E, Bray E, Lyapichev K, Ramos J, Krishnan V, Lai-Hsu C, Park KK, Tsoulfas P, et al. Perivascular Fibroblasts Form the Fibrotic Scar after Contusive Spinal Cord Injury. *Journal of Neuroscience*. 2013;33:13882–13887.
16. Zhou T, Zheng Y, Sun L, Badea SR, Jin Y, Liu Y, Rolfe AJ, Sun H, Wang X, Cheng Z, et al. Microvascular endothelial cells engulf myelin debris and promote macrophage recruitment and fibrosis after neural injury. *Nat Neurosci*. 2019;22:421–435.

17. Johnson CO, Nguyen M, Roth GA, Nichols E, Alam T, Abate D, Abd-Allah F, Abdelalim A, Abraha HN, Abu-Rmeileh NM, et al. Global, regional, and national burden of stroke, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet Neurology*. 2019;18:439–458.

## Figure Legends


**Figure 1.** The main constituent of chronic ICH lesions after hematoma resolution in a collagenase-induced rat model is collagen 3. Each row corresponds to a representative lesion in the striatum at 28 days (n=6), 42 days (n=6), and 73-85 days (n=6) post-ICH. Note that collagen 3 is seen in both thin, compacted lesions (42 days) as well as larger, less dense lesions (28 and 73-85 days). GFAP is predominantly observed at the lesion border. The white arrows indicate examples of structures of interest, and the white circles indicate examples of autofluorescence from hemosiderin. H&E scale bars (black) indicate 50µm and fluorescent image scale bars (white) indicate 100µm. For additional details on the lesion location, see Figure IX in the Data Supplement.

**Figure 2.** The lesion in chronic human ICH consists primarily of collagen 3. Each column corresponds to the indicated patient number (for additional case and location details, see the Table and Figure V in the Data Supplement). Each row is labeled by its corresponding marker. The top row provides an overview of the three chronic ICH lesions at low magnification followed by high magnification images that correspond to the dashed boxes. Elastic Van Gieson (eVG) staining at high magnification highlights (in pink) the extensive collagen found inside the lesions. Additional staining with collagen 3 and 4 reveals that the lesion consists primarily of collagen 3. All unlabeled scale bars after the first row indicate 100µm.



**Figure 1.** The main constituent of chronic ICH lesions after hematoma resolution in a collagenase-induced rat model is collagen 3. Each row corresponds to a representative lesion in the striatum at 28 days (n=6), 42 days (n=6), and 73-85 days (n=6) post-ICH. Note that collagen 3 is seen in both thin, compacted lesions (42 days) as well as larger, less dense lesions (28 and 73-85 days). GFAP is predominantly observed at the lesion border. The white arrows indicate examples of structures of interest, and the white circles indicate examples of autofluorescence from hemosiderin. H&E scale bars (black) indicate 50 $\mu$ m and fluorescent image scale bars (white) indicate 100 $\mu$ m. For additional details on the lesion location, see Figure S9 in the supplement.







**Figure 2.** The lesion in chronic human ICH consists primarily of collagen 3. Each column corresponds to the indicated patient number (for additional case and location details, see the Table and Figure S5 in the supplement). Each row is labeled by its corresponding marker. The top row provides an overview of the three chronic ICH lesions at low magnification followed by high magnification images that correspond to the dashed boxes. Van Gieson (eVG) staining at high magnification highlights (in pink) the extensive collagen found inside the lesions. Additional staining with collagen 3 and 4 reveals that the lesion consists primarily of collagen 3. All unlabeled scale bars after the first row indicate 100µm.